

**Baryon resonances in the mean field approach
and
a simple explanation of the Θ^+ pentaquark**

Dmitri Diakonov^{1,2}

¹*Petersburg Nuclear Physics Institute, Russia*

²*Institut für Theoretische Physik II,
Ruhr-Universität, Bochum, Germany*

(Dated: December 17, 2008)

Abstract

We suggest to classify baryon resonances as single-quark states in a mean field, and/or as its collective excitations. Identifying the Roper resonance $N(1440, \frac{1}{2}^+)$, the nucleon resonance $N(1535, \frac{1}{2}^-)$, and the singlet hyperon $\Lambda(1405, \frac{1}{2}^-)$ as single-quark excitations, we find that there must be an exotic $S = +1$ baryon resonance Θ^+ (the “pentaquark”) with a mass about $1440 + 1535 - 1405 = 1570$ MeV and spin-parity $\frac{1}{2}^+$. We argue that Θ^+ is an analog of the Gamov–Teller excitation long known in nuclear physics.

It was argued 30 years ago by Witten [1] that if the number of colors N_c is large, the N_c quarks constituting a baryon can be viewed as moving in a mean field whose fluctuations are suppressed as $1/N_c$. Whether $N_c=3$ of the real world is large enough for the mean field in baryons to be a working notion is a question to which there is no general answer: it depends on how large are $1/N_c, 1/N_c^2, \dots$ corrections to a particular baryon observable. However, experience in hadron physics tells us that usually the relations between observables found in the large- N_c limit are in satisfactory agreement with reality, unless there are special reasons to expect large $1/N_c$ corrections [1, 2]. In any case, it is helpful to understand how baryons are constructed in the large- N_c limit, before corrections are considered.

The mean field can, in principle, have components with various quantum numbers $J^{PC} = 0^{++}$ (scalar), 0^{-+} (pseudoscalar), 1^{--} (vector), 1^{++} (axial), etc. Since the mean field is created by N_c quarks of the u, d, s flavor it is also characterized by the flavor quantum numbers like isospin T and hypercharge Y . We do not consider baryons with heavy quarks here.

One expects that the mean field inside heavy ground-state baryons has the maximal possible, that is spherical symmetry. There is no problem in writing a spherical-symmetric scalar, flavor-singlet field as $\sigma(\mathbf{x}) = P_1(r)$ where $r = |\mathbf{x}|$ is a distance from the center of a baryon, however we immediately run into a problem of how to write the *Ansatz* for, say, the mean pseudoscalar field. Being pseudoscalar it has to be odd in $\mathbf{x}$. The minimal extension of spherical symmetry is then the “hedgehog” *Ansatz* “marrying” the isotopic and space axes:

$$\pi^a(\mathbf{x}) = n^a P_2(r), \quad n^a = \frac{x^a}{r}, \quad a = 1, 2, 3; \quad \pi^{4,5,6,7,8}(\mathbf{x}) = 0. \quad (1)$$

This *Ansatz* breaks spontaneously the symmetry under independent space and isospin rotations, and only a simultaneous rotation in both spaces remains a symmetry. At the same time it breaks the $SU(3)$ flavor symmetry. One may argue that the $SU(3)$ symmetry is explicitly broken from the start by $m_s \gg m_{u,d}$, however one can as well consider the strange quark mass m_s as a small perturbation [3]. In the chiral limit, $m_s \rightarrow 0$, the *Ansatz* (1) breaks spontaneously the $SU(3)$ symmetry: the first three component of the pseudoscalar octet are privileged. Full symmetry is restored when one rotates the asymmetric mean field in flavor and ordinary spaces: that produces many baryon states with definite quantum numbers.

The rôle of a small m_s is in fact the same as of the infinitesimal magnetic field in materials with a spontaneous magnetization: it establishes the preferred direction of magnetization.

In our case, the *Ansatz* (1) is privileged since $m_s \neq 0$, regardless of whether it is considered sizable or infinitesimal.

If in a baryon there are mean vector fields with the quantum numbers of ω and ϕ (and there are no *a priori* reasons why they should be absent), the *Ansätze* for those fields in correspondence with Eq. (1) are $\omega_0, \phi_0(\mathbf{x}) = P_{3,4}(r)$; $\omega_i, \phi_i(\mathbf{x}) = n_i P_{5,6}(r)$. The *Ansatz* for the axial a_1 field is $A_0^a = n^a P_7(r)$, $A_i^a = \epsilon_{aij} n_j P_8(r) + \delta_i^a P_9(r) + n^a n_i P_{10}(r)$, and so on.

In the mean field approximation, justified at large N_c , one looks for the solutions of the Dirac equation for single quark states in the background mean field. The Dirac Hamiltonian for quarks is, schematically,

$$H = \gamma^0 \left(i\gamma^i \partial_i + \sigma(\mathbf{x}) + i\gamma^5 \pi(\mathbf{x}) + \gamma^\mu V_\mu(\mathbf{x}) + \gamma^\mu \gamma^5 A_\mu(\mathbf{x}) + \dots \right) \quad (2)$$

In fact, the one-particle Dirac Hamiltonian is most probably nonlocal and momentum-dependent (as it would follow *e.g.* from Fiertz-transforming and then bosonizing color quark interactions), therefore Eq. (2) is a symbolic presentation. However, several important statements can be made on general grounds:

- Given the *Ansatz* for the mean fields σ, π, V, A , the Hamiltonian (2) actually splits into two: one for s quarks (H_s) and the other for u, d quarks (H_{ud}). The former commutes with the angular momentum of s quarks, $\mathbf{J} = \mathbf{L} + \mathbf{S}$, and with the inversion of spatial axes, hence all energy levels of s quarks are characterized by half-integer J^P and are $(2J + 1)$ -fold degenerate. The latter commutes only with the ‘grand spin’ $\mathbf{K} = \mathbf{T} + \mathbf{J}$ and with inversion, hence the u, d quark levels have definite integer K^P and are $(2K + 1)$ -fold degenerate. The energy levels for u, d quarks on the one hand and for s quarks on the other are completely different, even in the chiral limit $m_s \rightarrow 0$
- The Hamiltonian (2) has mixed symmetry with respect to time reversal, therefore the one-particle spectra for s and u, d quarks are generally not symmetric under the change $E \rightarrow -E$
- All energy levels, both positive and negative, are probably discrete owing to confinement. Indeed, a continuous spectrum would correspond to a situation when quarks are free at large distances from the center, which contradicts confinement. [One can mimic or model confinement for example by imposing the condition that the effective quark mass $\sigma(\mathbf{x})$ grows at infinity.]

- According to the Dirac theory, all *negative-energy* levels, both for s and u, d quarks, have to be fully occupied, corresponding to the vacuum. It means that there must be exactly N_c quarks antisymmetric in color occupying all (degenerate) levels with J_3 from $-J$ to J , or K_3 from $-K$ to K ; they form closed shells that do not carry quantum numbers
- Filling in the lowest level with $E > 0$ by N_c quarks makes a baryon.

We shall suppose that the energy levels with minimal $|E|$ are $K^P = 0^+$ for u, d quarks and $J^P = \frac{1}{2}^+$ for s quarks, because it corresponds to the maximal-symmetry wave functions as it is usual for the Dirac equation in smooth external fields. *A priori* the signs of E for those solutions can be any, however experimentally the lowest baryon is a nucleon and not the Ω^- hyperon. It means, that the $\frac{1}{2}^+$ level for s quarks is probably the nearest to $E = 0$ but remains on the negative side. It belongs to the vacuum sector and has to be filled in together with all the rest negative-energy levels. On the contrary, the 0^+ level for u, d quarks must be the lowest at $E > 0$. Filling it in with N_c quarks antisymmetric in color, adds N_c u, d quarks to the vacuum, and makes the nucleon, see Fig. 1. We do not know yet how much higher is the highest filled u, d shell than the highest filled s -quark shell but shall determine in shortly from experiment (at $N_c = 3$).




FIG. 1: Filled quark levels for the ground-state baryon $N(940, \frac{1}{2}^+)$. The two lightest baryon multiplets $(8, \frac{1}{2}^+)$ and $(10, \frac{3}{2}^+)$ are rotational excitations of the same filling scheme.

The true quantum numbers of the lightest baryons are determined from the quantization of the rotations of the mean field since the *Ansatz* discussed above spontaneously breaks symmetry under rotations in ordinary and flavor spaces. However, a simultaneous rotation in ordinary and isospin spaces remains a symmetry. Therefore, if one limits oneself to non-strange baryons, the quantization of rotations produces states with $J = T = \frac{1}{2}$ (for any odd N_c), that is the nucleon, and $J = T = \frac{3}{2}$, the Δ resonance [4]. If m_s is treated as a

perturbation [3], one has to extend this to $SU(3)$ flavor rotation. Its quantization gives specific $SU(3)$ multiplets that reduce at $N_c = 3$ to the octet with spin $\frac{1}{2}$ and the decuplet with spin $\frac{3}{2}$, see *e.g.* [5]. Witten's quantization condition $Y' = \frac{N_c}{3}$ [4] follows trivially from the fact that there are N_c u, d valence quarks each with the hypercharge $\frac{1}{3}$ [6]. Therefore, the ground state shown in Fig. 1 entails in fact 56 rotational states. It is the same **56**-plet as the ground shell of the nonrelativistic quark model but its interpretation is different. The splitting between the centers of the multiplets $(\mathbf{8}, \frac{1}{2}^+)$ and $(\mathbf{10}, \frac{3}{2}^+)$ is $\mathcal{O}(1/N_c)$, and the splittings inside multiplets can be determined as a linear perturbation in m_s [6].

The picture has a similarity in nuclear physics where at large atomic numbers A , Z protons and $A-Z$ neutrons are considered in different self-consistent mean fields and have a different system of one-particle levels. One fills proton and neutron levels separately up to the common Fermi surface. Contrary to the quark case, the negative-energy levels for nucleons can be neglected because of the large nucleon mass. There are collective excitations of heavy nuclei, *e.g.* rotation whose energy scale is $\mathcal{O}(1/A)$ (in the baryon case it scales as $1/N_c$). However, there are also one-particle and particle-hole excitations that are of the order of unity in A . Similarly, one should expect $\mathcal{O}(N_c^0)$, that is large, one-particle and particle-hole excitations. Let us try to identify them.

The lowest baryon resonance beyond the rotational excitations of the ground state is the singlet $\Lambda(1405, \frac{1}{2}^-)$. Apparently, it can be obtained only as an excitation of the s quark, and its quantum numbers must be $J^P = \frac{1}{2}^-$. The resonance $\Lambda(1405, \frac{1}{2}^-)$ is excited if one of the u, d quarks from the valence 0^+ level jumps, under the action of a $S = -1$ force, to the first excited state for s quarks, see Fig. 2. It is predominantly a 3-quark state (at $N_c = 3$). This excited state generates a rotational band of $SU(3)$ multiplets of its own, but we do not consider them here.




FIG. 2: $\Lambda(1405, \frac{1}{2}^-)$




FIG. 3: $N(1535, \frac{1}{2}^-)$

If there is a $\frac{1}{2}^-$ level for s quarks, it can be excited also by an s quark jumping from its highest filled level $\frac{1}{2}^+$, see Fig. 3. This is a particle-hole excitation which does not change the nucleon quantum numbers (except for parity), as we are just adding an $s\bar{s}$ pair to the valence u, d level that determines the quantum numbers. We therefore identify this excitation with $N(1535, \frac{1}{2}^-)$. We see that at $N_c=3$ it is predominantly a pentaquark state $u(d)uds\bar{s}$. That explains its large branching ratio in the ηN decay [7], a long-time mystery. We also see that, since the highest filled level for s quarks is lower than the highest filled level for u, d quarks, $N(1535, \frac{1}{2}^-)$ must be *heavier* than $\Lambda(1405, \frac{1}{2}^-)$: the opposite prediction of the nonrelativistic quark model has been always of some concern. We stress that in our picture the existence of an unusual nucleon resonance $N(1535, \frac{1}{2}^-)$ is a *consequence* of the $\Lambda(1405, \frac{1}{2}^-)$ existence. The transition shown in Fig. 3 also entails its own rotational band. Subtracting $1535 - 1405 = 130$, we find that the $\frac{1}{2}^+$ s -quark level is approximately 130 MeV lower in energy than the valence 0^+ level for u, d quarks.

There is also a low-lying Roper resonance $N(1440, \frac{1}{2}^+)$. It requires that there is an excited one-particle u, d state with $K^P = 0^+$, see Fig. 4. Just as the ground state nucleon, it is part of the excited $(\mathbf{8}', \frac{1}{2}^+)$ and $(\mathbf{10}', \frac{3}{2}^+)$ split as $1/N_c$. In fact the first excited state could be also $K^P = 1^+, 2^+$ which would generate more $SU(3)$ multiplets including one with the Roper resonance; $K^P = 0^+$ is a minimal hypothesis. The identification of the nature of the Roper resonance solves another problem of the nonrelativistic model where $N(1440, \frac{1}{2}^+)$ must be heavier than $N(1535, \frac{1}{2}^-)$. In our approach they are simply unrelated.




FIG. 4: $N(1440, \frac{1}{2}^+)$




FIG. 5: $\Theta^+(\frac{1}{2}^+)$

We now come to the crucial point: Given that there is an unoccupied level for u, d quarks, one can put there an s quark as well, taking it from one of the filled s -quark shells. The minimal-energy excitation is from the highest occupied shell for s quarks, to the lowest unoccupied level for u, d quark, that is to the would-be Roper level, see Fig. 5. It is a particle-hole excitation with the valence level left untouched, its quantum numbers being

apparently $S = +1$, $T = 0$, $J^P = \frac{1}{2}^+$. At $N_c = 3$ this excitation is a pentaquark state $uudd\bar{s}$, precisely the exotic Θ^+ baryon predicted in Ref. [8] from other considerations. The quantization of its rotations in flavor and ordinary spaces produces the antidecuplet $(\overline{\mathbf{10}}, \frac{1}{2}^+)$ and higher multiplets $(\mathbf{27}, \frac{3}{2}^+)$ and $(\mathbf{27}, \frac{1}{2}^+)$ [9].

Since the relative position of all four levels involved are already known from the masses of the well-established resonances $N(1440, \frac{1}{2}^+)$, $N(1535, \frac{1}{2}^-)$ and $\Lambda(1405, \frac{1}{2}^-)$, it is a matter of trivial arithmetics to find the energy difference between the s -quark shell and the first excited u, d -quark level. We obtain an estimate for the Θ^+ mass: $m_\Theta \approx 1440 + 1535 - 1405 = 1570$ MeV. Of course, one should not understand the number literally. First, the masses of the resonances exploited here are known with an uncertainty of a few tens MeV. Second, the masses of physical resonances are not fixed exactly by the one-particle levels but get $\mathcal{O}(1/N_c)$ and $\mathcal{O}(m_s)$ corrections (which in principle are calculable). Nevertheless, the most interesting prediction that the exotic pentaquark is a consequence of the three well-known resonances and must be light, is an unambiguous feature of the picture.

In nuclear physics, the charge-exchange excitations generated by the axial current $j_{\mu 5}^\pm$, when a neutron from the last occupied shell is sent to an unoccupied proton level or *vice versa* are known as Gamov–Teller transitions [10]; they have been extensively studied both experimentally and theoretically. Thus our interpretation of the Θ^+ is that it is a Gamov–Teller-type resonance long known in nuclear physics.

However, the calculation of the Θ^+ width should be different from that in nuclear physics. For the Gamov–Teller transition in a nuclei, it is sufficient to calculate the matrix element between the initial and final states of the transition axial current. To make it more accurate, one can take into account a correction from the admixture of particle-hole states to the one-particle states. This approximation is often called the RPA; it goes beyond the mean field.

In the baryon case, even a one-particle state of the leading mean-field approximation, shown in Fig. 1, is in fact a Fock state with additional quark-antiquark pairs. These arise when one decomposes the filled negative-energy levels in the plane wave basis [11, 12]. Therefore, unlike its nuclear physics counterpart, the Θ^+ decay amplitude has, already in the mean field approximation, two contributions: one is from the conventional “fall-apart” process whereas the other is the “5-to-5” transition of the Θ^+ to the 5-quark component of the nucleon [5, 13]. The two amplitudes are separately not Lorentz invariant – only their sum is. In the lab frame there is a tendency for the two amplitudes to cancel each

other [14]. In the infinite momentum frame, however, the “fall-apart” amplitude (the simple one) is zero in the chiral limit, and only the “5-to-5” amplitude survives. The one-particle Hamiltonian (2) is covariant, such that there is no problem in transforming the mean field to an infinite momentum frame, which is the shortest way to evaluate the Θ^+ width. [It may seem somewhat unusual but we do not have much experience from the past in computing pentaquarks widths!] The program has been carried out in the Chiral Quark Soliton Model with the result $\Gamma_\Theta \sim 1 \text{ MeV}$ with no parameter fitting [12, 15].

It should be stressed that the small width of Θ^+ has no relation to and no influence from the large width of the Roper resonance. When Θ^+ decays, the valence level (to which the Roper resonance decays) is Pauli-blocked, and the fact that an ordinary “3-to-3” decay width of the Roper resonance is large does not effect the width of the Θ^+ which is very narrow owing to physics completely different from that governing the Roper decay.

There can be additional one-particle and/or particle-hole excitations, however just the two excited levels suggested here (0^+ for u, d quarks and $\frac{1}{2}^-$ for s quarks) are sufficient to explain the majority of baryon resonances up to 2 GeV, if the rotational states generated by each of the excitations are taken into account. A detailed study of the ensuing baryon spectrum will be published separately. For high-spin resonances (actually for $J \geq N_c$) it may become energetically favorable to depart from a spherically-symmetric mean field.

I am grateful to L. Frankfurt, K. Goeke, A. Hosaka, M. Polyakov and especially to V. Petrov for helpful discussions. I thank German Research Society (DFG) for Mercator Professorship at Bochum. This work has been partly supported by Russian Government grants RFBR-06-02-16786 and RSGSS-3628.2008.2.

-
- [1] E. Witten, Nucl. Phys. **B160**, 57 (1979)
 - [2] A. Manohar, “Hadrons in the $1/N$ expansion”, in: At the frontier of particle physics, M. Shifman (ed.), World Scientific, vol. 1, 507 (2002); Large N_c QCD 2004, Proceedings Int. Workshop, Trento, July 5-11, 2004, J.L. Goity, R.F. Lebed, A. Pich, C.L. Schat, N.N.Scoccola (eds.), World Scientific (2005) 307 p.; T. Cohen, Nucl. Phys. **A755**, 40 (2005), arXiv:hep-ph/0501090
 - [3] The lowest $SU(3)$ baryon multiplets satisfy Gell-Mann–Okubo relations with a high accuracy,

meaning that the splittings inside the multiplets are linear in m_s , actually $\mathcal{O}(m_s N_c)$. The splitting between the centers of the octet and of the decuplet is $\mathcal{O}(\Lambda/N_c)$ where Λ is the QCD scale parameter. Numerically, the former quantity is about 140 MeV whereas the latter is $1382 - 1152 = 230$ MeV, slightly larger. It means that in baryon physics one can indeed treat m_s as a very small quantity: $m_s = \mathcal{O}(\Lambda/N_c^2)$ or even less.

- [4] E. Witten, Nucl. Phys. **B223**, 433 (1983)
- [5] D. Diakonov and V. Petrov, to be published in *Multifaceted Skyrmions*, G. Brown and M. Rho, eds., World Scientific, arXiv:0812.1212 [hep-ph]
- [6] A. Blotz, D. Diakonov, K. Goeke, N.W. Park, V. Petrov and P. Pobylitsa, Nucl. Phys. **A355**, 765 (1993)
- [7] B.S. Zou, Eur. Phys. J. **A35**, 325 (2008), arXiv:0711.4860 [nucl-th]
- [8] D. Diakonov, V. Petrov and M. Polyakov, Zeit. Phys. **A359**, 305 (1997), arXiv:hep-ph/9703373
- [9] D. Diakonov and V. Petrov, Phys. Rev **D69**, 056002 (2004), arXiv:hep-ph/0309203
- [10] A. Bohr and B. Mottelson, Nuclear structure, W. A. Benjamin, NY (1998)
- [11] V. Petrov and M. Polyakov, arXiv:hep-ph/0307077
- [12] D. Diakonov and V. Petrov, Phys. Rev. **D72**, 074009 (2005), arXiv:hep-ph/0505201
- [13] D. Diakonov, Prog. Theor. Phys. Suppl. **168**, 1 (2007), and hep-ph/0610166
- [14] A. Hosaka, private communication
- [15] C. Lorcé, Phys. Rev. **D74**, 054019 (2006), arXiv:hep-ph/0603231; T. Ledwig, H.-C. Kim and K. Goeke, Phys. Rev. **D78**, 054005 (2008), arXiv:0805.4063